

Cancellation of $1/m_Q$ Corrections to the Inclusive Decay Width of a Heavy Quark *

A. Sinkovics

The Randall Laboratory of Physics

University of Michigan

Ann Arbor, MI 48109-1120

I. INTRODUCTION

This talk is a short summary of a work done in collaboration with Ratindranath Akhoury and Valentine Zakharov [1], a study of power corrections to the inclusive semileptonic decay of a heavy quark. The phenomenological motivation of this work is that the semileptonic inclusive decay can provide information on the CKM matrix elements. Power corrections may be large and knowledge about them is important for the precision extraction of the CKM parameters.

Inclusive decays of hadrons containing a single heavy quark (mass m_b) have been studied [2] within the operator product expansion (OPE) which gives an expansion of the decay width in terms of a small parameter Λ/m_b :

$$\Gamma_{incl} = \Gamma_{incl}^{pert}(1 + a_1\Lambda/m_b + a_2(\Lambda/m_b)^2 + \dots). \quad (1)$$

In the above, Γ_{incl}^{pert} is the decay width of the heavy quark including all perturbative corrections (the parton model result). Since there are no local gauge invariant operators of the appropriate dimension, an important conclusion [2] is that $a_1 = 0$, and the higher order power corrections can be classified in terms of the matrix elements of various operators.

Power corrections may be studied with the method of renormalons [3] within perturbation theory. The basic idea of the renormalon method is that one can obtain information about power corrections by looking at a class of diagrams which give factorial divergence in large orders of perturbation theory. The power corrections are seen to arise from regions of low momenta. This leads us to investigate the infrared sensitivity of inclusive decays, and we can replace the QCD scale Λ , which is a non-perturbative parameter, by an infrared cutoff parameter λ defined within perturbation theory. In the case of QED, for example, it could be a (fictitious) photon mass. Then the inclusive rate (1) may be represented as :

$$\Gamma_{incl} = \Gamma_{incl}^{pert}(1 + b_1\lambda/m_b + b_2(\lambda/m_b)^2 \ln \lambda + \dots). \quad (2)$$

Note that the Landau conditions for the singularities of Feynman diagrams tell us that the infrared sensitive contributions arise only as terms which are non-analytic in λ^2 .

An advantage of the OPE is the generality of its applicability to non-perturbative contributions. On the other hand, the conditions for the validity of the OPE can be checked by explicit perturbative calculations. As mentioned before, since no dimension 4 local gauge

*Talk given at the 36th International School of Subnuclear Physics, Erice

invariant operator in HQET, the coefficient of the first, linear correction $b_1 = 0$. Explicit one-loop order calculations [4] however have found linear correction to the decay width ($b_1 \neq 0$). The reason is that the pole mass is infrared sensitive, so on-shell renormalization must be avoided, and one has to use a short-distance mass instead. The decay width has no linear correction only if expressed not in terms of the pole mass m_b but in terms of a short distance or a $\bar{M}S$ mass. In this way one avoids an otherwise large (possibly non-universal) correction of order $1/m_b$, but at the expense of introducing a renormalization scale (μ) dependence in the total rate through the use of the running mass.

We extended the above argument for the inclusive semileptonic decay of a heavy quark in QCD to the second loop order when the non-abelian nature of the interactions are first apparent. That is we show the cancellations of all terms linear in the infrared cutoff in the decay width when the latter is expressed in terms of the short distance mass. The possible sources of infrared sensitive terms are the mass and wave function renormalization diagrams of the initial heavy quark and the bremsstrahlung diagrams. The cancellations proceed in a different manner for the two cases because whereas the wave function renormalization is multiplicative, the on-mass shell renormalization is additive.

II. CONNECTION TO THE KINOSHITA-LEE-NAUENBERG THEOREM

Certain questions immediately come to mind concerning the generality of the result. Is there a general principle behind this cancellation?, and is it true to all orders in perturbation theory, even for the non-abelian case? In fact there are infrared safe observables in QCD that do receive power corrections that are proportional to $1/M$ where M is a large mass scale. Examples are provided by the event shape variables in e^+e^- annihilation [5] for which there does not exist an operator product expansion. It has been argued in ref. [6] that inclusive enough observables do not receive $1/M$ power corrections whereas the more exclusive ones that assume some precision measurement do. Below we will apply this principle to the inclusive decay of the heavy quark, and argue that the KLN [7] theorem is behind the cancellation of the terms proportional to $1/m_b$.

The KLN theorem states the cancellation of infrared divergences in the transition probability when it is summed over an appropriate degenerate set of initial and final states. In a recent publication [8] it was argued that in the KLN sum not only the leading infrared sensitive piece proportional to $\ln\lambda$ but also the next, subleading $\sim \lambda$ term is cancelled, i.e.,

$$\sum_{i,f} |S_{i \rightarrow f}|^2 \sim 0 \cdot \ln\lambda + 0 \cdot \lambda + \text{terms independent of } \lambda + O(\lambda^2 \ln\lambda). \quad (3)$$

The transition probability summed over both initial and final states cannot be related to a physical inclusive cross section. However we argue that for heavy quark decay the initial state is trivial and hence from the above the KLN theorem guarantees the cancellation not only of the $\ln\lambda$ terms but also the subleading linear pieces. According to the uncertainty principle, the total energy can be measured to any accuracy provided the measurement time is long enough. In particular, the uncertainty ΔE in the total energy can be made smaller than an infrared cutoff:

$$\Delta E \ll \lambda \quad (4)$$

Now consider a decaying charged (or colored) particle in its rest frame. It is obvious that this state is not degenerate with any other since there is a gap between the mass of the charged particle and the energies in the continuum for any $\lambda \neq 0$. Thus the summation over the initial states in the KLN theorem is redundant for this case. Hence we conclude that

$$\Gamma_{incl} = \Gamma_{incl}^{pert} \left(1 + O(\lambda^2 \ln \lambda)\right). \quad (5)$$

It should be emphasized that this argument guarantees the above equality to all orders in perturbation theory. However it is very important for the KLN theorem to be valid that the renormalization procedure does not introduce any infrared sensitivity. Thus since the pole mass is infrared sensitive, on shell mass renormalization must be avoided in favour of the $\bar{M}S$ scheme. We would also like to point out that the same argument can be applied not just to the semileptonic decay but to other inclusive decays like for example the radiative one, $B \rightarrow X_s \gamma$.

As mentioned before, we verify the above argument by an explicit second loop order calculation: we show the cancellation of terms linearly proportional to an infrared cutoff λ . Our method, discussed in section III, is to introduce local effective vertices for the diagrams with radiation from the final state quark to linear accuracy.

III. EFFECTIVE VERTICES

We consider the totally inclusive semileptonic decay of a heavy quark and examine the possible terms linearly proportional to an infrared cutoff. Such terms can arise from the soft gluon contributions to the diagrams for the mass and wave function renormalizations of the heavy quark and from the bremsstrahlung diagrams.

The decay rate is proportional to the imaginary part of the forward amplitude shown in Fig. 1. The hard amplitude involving the final state quark is denoted by the shaded blob which also includes the lepton loop. The soft gluon interactions responsible for the infrared sensitivity dress the hard amplitude, connecting to it and to the heavy quark. We use the fact that for the soft gluon of momenta $k \ll m_b$, we may perform the sum over cuts implicit in taking the imaginary part, for just the hard part of the amplitude leaving the soft gluon lines uncut. Thus we sum over the cuts of the hard amplitude only and leave the soft gluon propagators uncut in the diagrams for the forward scattering amplitude. One can show that up to linear accuracy, after summing over the cuts of the hard part, and integrating over its phase space and that of the leptons, the interaction of soft gluon radiation with the hard part may be replaced by effective local vertices.

In the parton model, the total width for the semileptonic decay is to lowest order:

$$\Gamma_0 = \frac{G_F^2 |V_{ub}|^2}{192\pi^3} m_b^5 \quad (6)$$

We may represent this by an effective local vertex, obtained by performing the phase space integration over the leptons and the massless quark. Such an effective vertex may be written as:

$$C \bar{u} \not{p}^5 u, \quad C = \frac{G_F^2 |V_{ub}|^2}{192\pi^3}. \quad (7)$$

where u is the heavy quark spinor. This vertex shown on Fig. 2a) and will be referred to as the leading effective vertex.

Similarly, effective vertices can be derived for the absorption of one and two gluons from the final state quark. (Figure 2b), 2c)). These effective vertices may be obtained from the leading one by gauge invariance, by means of the replacement $p_\mu \rightarrow (p_\mu - gT^a A_\mu^a)$. In the following we study the infrared contribution of the effective vertex diagrams and show the cancellation of infrared sensitive pieces linear in an infrared cutoff.

All the infrared cancellations are shown strictly algebraically at the level of the corresponding Feynman integrands. Power counting is used to isolate and to show the cancellations of the possibly infrared sensitive contributions upto linear accuracy. In this way we avoid using any particular IR cutoff, which is crucial for the case of non-abelian theory.

IV. CANCELLATION OF INFRARED SENSITIVE TERMS UPTO LINEAR ACCURACY FROM THE WAVE FUNCTION RENORMALIZATION DIAGRAMS

In this section we will show that the infrared sensitive terms upto linear accuracy are cancelled between the wave function renormalization contributions from diagrams with self energy insertions and parts of bremsstrahlung diagrams. Since the wave function renormalization is multiplicative, all its contributions factorize from the hard amplitude. The corresponding bremsstrahlung diagrams that cancel the logarithmic and linear dependence on the IR cutoff of the wave function renormalization diagrams may be obtained by certain Ward-like identities. In this talk I explicitly show our method on the the one-loop order cancellation only. The procedure can be generalized to second loop order, which is the actual new result.

We will denote by $-i\Sigma(p)$ the sum of all 2 point one particle irreducible graphs. Then the full fermion propagator is:

$$S'_F(p) = \frac{i}{\not{p} - m_b - \Sigma(p)} \quad (8)$$

Near the mass shell we have

$$\Sigma(p) = \delta m(m_b, \Delta m) - (Z^{-1} - 1)(\not{p} - m_b) + O((\not{p} - m_b)^2) \quad (9)$$

where Δm denotes the mass counterterm:

$$\delta m(m_b, \Delta m) |_{\not{p}=m_b} = 0. \quad (10)$$

Thus m_b denotes the pole or the physical mass:

$$\lim_{p \rightarrow m_b} S'_F(p) = \frac{i}{\not{p} - m_b} Z \quad (11)$$

From the LSZ reduction formula, each external line in the S-matrix element gets a factor $Z^{1/2}$ and we are interested in its perturbative expansion. Thus we expand,

$$Z^{1/2} = 1 - \frac{1}{2}(Z^{-1} - 1) + \frac{3}{8}(Z^{-1} - 1)^2 + \dots \quad (12)$$

and $Z^{-1} - 1$ may be computed using:

$$Z^{-1} - 1 = - \frac{\bar{u} \frac{\partial}{\partial p_\nu} \Sigma|_{\not{p}=m_b} u}{\bar{u} \gamma^\nu u} \quad (13)$$

In the rest frame of the heavy quark, only $\nu = 0$ contributes.

Consider the one-loop order wave function renormalization and the corresponding bremsstrahlung diagrams shown on Fig. 3. The wave function renormalization parts of both graph(a) and graph(b) are the hard part times a factor of $Z^{1/2}$. Thus we expand it using (12) keeping only the first two terms and computing the perturbative contribution to order α of $Z^{-1} - 1$ from (13). Next using the identity

$$\frac{\partial}{\partial p^\nu} \frac{i}{\not{p} - \not{k} - m_b} = \frac{i}{\not{p} - \not{k} - m_b} i\gamma_\nu \frac{i}{\not{p} - \not{k} - m_b} \quad (14)$$

we obtain the Ward-like identity shown in Fig. 4, where the dotted line denotes the insertion of a zero momentum vertex γ_ν . Going to the rest frame of the heavy quark and using Eqs. (12), (13) we immediately see

$$3c)|_{wave\ function} = (Z^{-1} - 1)g^2 C m_b^5 \bar{u}u = -[3a) + 3b)]. \quad (15)$$

Here, and henceforth, unless otherwise specified, the equality sign is for terms that are logarithmically and linearly infrared sensitive. This is the desired result, which does not involve using an explicit infrared cutoff.

The cancellation of the wave function parts at two loops follows the same general method except that now we have more diagrams. The 2-loop one particle irreducible diagrams can be divided into 5 groups according to the 5 pieces of the order g^4 self-energy corrections (Fig. 6). These groups are essentially the different color groups, with the exception that the first and third group have the same color structure. The cancellation of logarithmic and linear terms in the infrared cutoff takes place within each color group separately. As in the one-loop case, the cancellation is obtained by use of Ward-like identities, valid to linear accuracy. Figures 7 and 8 show one color group of diagrams and the corresponding Ward-like identity as an example. As before, the infrared sensitive pieces are identified by power counting (no specific cutoff is used) and the cancellation is algebraic at the level of integrands.

At the two-loop level 1-particle reducible diagrams also contribute to the wave function renormalization. The contribution of the diagrams can be simply calculated using the perturbative expansion of $Z^{1/2}$ (Eq. 12) multiplying each external line in the S-matrix element. A straightforward calculation shows that the wave function renormalization pieces from the two-loop order 1-particle reducible diagrams cancel.

V. CANCELLATION OF THE INFRARED SENSITIVE TERMS FROM MASS RENORMALIZATION

Let us now consider the cancellation of the leading infrared sensitive piece (the one that is linearly divergent in the infrared) arising from the diagrams involving the mass renormalization of the heavy quark. While the wave function renormalization was multiplicative,

and so the hard part of diagrams factorized, the mass renormalization is additive, and the structure of hard vertex is important here. As for the wave function pieces, I discuss the cancellation in one loop order to exemplify our method, and indicate how to generalize the calculation for second loop order.

As discussed in the introduction, the pole mass of the heavy quark contains a long distance piece which is proportional linearly to an infrared cutoff, the first contribution starting at order α . By infrared power counting we can isolate the linearly infrared sensitive pieces of the self-energy diagrams as

$$\begin{aligned}\Sigma_1|_{\not{p}=m_b} &= a_1 g^2 + b_1 g^2 \lambda \\ \Sigma_{2i}|_{\not{p}=m_b} &= a_{2i} g^4 + b_{2i} g^4 \lambda \quad i = 1 \dots 5\end{aligned}\tag{16}$$

It is convenient to separate out this piece and define the short distance or running mass to one and second loop order respectively as

$$\begin{aligned}\bar{m} &= m_0 + a_1 g^2 \\ \bar{m} &= m_0 + a_1 g^2 + \sum_{i=1}^5 a_{2i} g^4.\end{aligned}\tag{17}$$

We should emphasize our notation at this point. Even though one may use a gluon mass as an infrared cutoff at the one loop level, this is not a gauge invariant procedure beyond it. Thus λ refers to some gauge invariant cutoff, which we need not specify. Terms that are linearly divergent in the infrared are identified by power counting and cancellation of such terms is shown at the level of the corresponding Feynman integrands.

The various diagrams contributing to this order are shown in Fig. 5. Fig. 5a) is the bremsstrahlung contribution. To find the leading infrared sensitive contribution, we expand the leading effective hard vertex to first order in the offshellness,

$$C(\not{p} - \not{k})^5 = C(\not{p} - \not{k} - m_b + m_b)^5 = C m_b^5 + 5C m_b^4 (\not{p} - \not{k} - m_b) + \dots\tag{18}$$

This is an expansion of the leading effective vertex in $(\not{p} - \not{k} - m_b)/m_b$ and it is understood that we keep such deviations from the mass shell in as much as they cancel the corresponding small denominators from the propagators. Then after this expansion, one of the fermion propagators in the integrand of the expression for Fig. 5 is cancelled (shown by a slash in the Figure) and we are left with an expression which resembles that for the self energy:

$$5a) = 5C(\bar{m})^4 b_1 g^2 \lambda \bar{u} u\tag{19}$$

Next we have the contributions from the bremsstrahlung from the final state quark which as discussed earlier is replaced by the effective vertices for the gluon emission. The two diagrams with single gluon effective vertices of Figs. 5b) and 5c) give:

$$5b) + 5c) = 2 \times C(-i5m_b^4) \bar{u} (-i\Sigma_1(p)) u = -2 \times 5C(\bar{m})^4 b_1 g^2 \lambda \bar{u} u.\tag{20}$$

Finally, in the lowest order diagram, Fig. 5d), we must express the leading effective vertex in terms of the short distance mass,

$$5d) = C m_b^5 = C(\bar{m} + b_1 g^2 \lambda)^5 = C \bar{m}^5 \bar{u} u + 5C(\bar{m})^4 b_1 g^2 \lambda \bar{u} u.\tag{21}$$

Adding together all the order g^2 terms we see the cancellation of the linearly infrared divergent pieces to this order when the decay rate is expressed in terms of the running mass $\bar{m}$.

As in the wave function renormalization, we group the two-loop 1 PI diagrams according to the five self-energy pieces shown in Fig. 6. These groups are now enlarged by new diagrams with the single and double gluon vertices (Fig. 2). Unlike in the wave function case, where each group of graphs cancelled separately, the cancellation of linear infrared sensitive pieces in the mass renormalization involves mixing between the different groups and mass shift terms from one loop graphs. To show the cancellation of linearly infrared sensitive pieces, we follow the same procedure as in the one-loop case: expand the effective vertices in the offshellness, keep terms that would be linearly proportional to an infrared cutoff, and show the cancellation algebraically. Two cancelling sets are shown on Figures 9, and 10. There are two additional sets of diagrams with abelian type of cancellation (the cancellation proceeds in the same way as the interaction was abelian). The second loop order calculation explicitly verifies that cancellation of linearly infrared pieces occurs if expressing the decay width in terms of the short distance (running) mass.

VI. SUMMARY AND CONCLUSIONS

We have shown by explicit perturbative calculations upto the second loop order that when the inclusive decay width of a heavy quark is expressed in terms of the short distance mass, there is complete accord with the operator product expansion and the leading power corrections are of order $1/m_b^2$.

As emphasized earlier, the above result is true not just for the semileptonic decay considered in this paper but for any inclusive decay with a single heavy particle initial state, like for example the radiative one. Such a cancellation was linked to the KLN theorem which had been suggested earlier [6] to be the general principle behind this for inclusive enough observables.

This work was supported in part by the US Department of Energy.

-
- [1] A. Sinkovics, R. Akhoury and V.I. Zakharov, hep-ph/9804401, to appear in *Phys. Rev.* **D**
 - [2] J. Chay, H. Georgi and B. Grinstein, *Phys. Lett.* **B247** (1990) 399; I. I. Bigi, N. G. Uraltsev, A. I. Vainshtein, *Phys. Lett.* **B293** (1992) 430; I. I. Bigi, M. Shifman, N. G. Uraltsev, A. I. Vainshtein, *Phys. Rev. Lett.* **71** (1993), 496; A. V. Manohar and M. B. Wise, *Phys. Rev.* **D 49** (1994), 1310.
 - [3] R. Akhoury and V.I. Zakharov, *Nucl. Phys. Proc. Supp.* **A 54** (1997) 217, hep-ph/9610492.
 - [4] M. Beneke and V. M. Braun, *Nucl. Phys.* **B426** (1994) 301; I. I. Bigi, M. Shifman, N. G. Uraltsev, A. I. Vainshtein, *Phys. Rev.* **D 50** (1994), 2234; M. Beneke, V. M. Braun and V. I. Zakharov, *Phys. Rev. Lett.* **73** (1993) 3058.
 - [5] B. R. Webber, *Phys. Lett.* **B339** (1994) 148; B. R. Webber and Yu. L. Dokshitzer, *Phys. Lett.* **B352** (1995) 415; R. Akhoury and V. I. Zakharov, *Phys. Lett.* **B357** (1995) 646.

- [6] R. Akhouri and V. I. Zakharov, *Phys. Rev. Lett.* **76** (1996) 2238.
 [7] T. Kinoshita *J. Math. Phys.* **3** (1962) 650; T. D. Lee and M. Nauenberg *Phys. Rev.* **133** (1964) 1549.
 [8] R. Akhouri, L. Stodolsky and V. I. Zakharov, *Nucl. Phys.* **B516** (1998) 317, hep-ph/9609638




FIG. 1. The forward scattering amplitude for the heavy quark decay whose imaginary part gives the decay width. Shaded blob denotes the hard part, while S denotes the soft gluon interactions.




FIG. 2. a) leading effective vertex, b) single gluon effective vertex , c) double gluon effective vertex




FIG. 3. Wave function renormalization and Bremsstrahlung diagrams to order g^2 .

$$-i \frac{\partial}{\partial p_\nu} \text{ (diagram with wavy line) } = \text{ (diagram with wavy line and index } \nu \text{) }$$

FIG. 4. One-loop order Ward-like identity




FIG. 5. Mass renormalization and the corresponding bremsstrahlung diagram to order g^2 . Figure a) depicts the mass renormalization piece of the graph. The propagator with slash is cancelled by the expansion of the leading effective vertex. The subscript Σ_1 of d) denotes the mass shift due to the order g^2 self-energy correction.

$$\begin{aligned}
 -i\Sigma = & \frac{\text{diagram 1}}{-i\Sigma_1} + \frac{\text{diagram 2}}{-i\Sigma_{21}} + \frac{\text{diagram 3}}{-i\Sigma_{22}} + \frac{\text{diagram 4}}{-i\Sigma_{23}} \\
 & + \frac{\text{diagram 5}}{-i\Sigma_{24}} + \frac{\text{diagram 6}}{-i\Sigma_{25}}
 \end{aligned}$$

FIG. 6. Self-energy corrections of order g^2 and g^4




FIG. 7. The second group of 2-loop diagrams corresponding to Σ_{22} .

$$-i \frac{\partial}{\partial p_\nu} \text{diagram 1} = \text{diagram 2} - \text{diagram 3} + \text{diagram 4}$$

FIG. 8. Ward-like identity to the second group of 2-loop diagrams




FIG. 9. Mass renormalization of order g^4 : group-1 and group-2. The linear infrared sensitive mass renormalization pieces of the leading effective vertex graphs cancel with the corresponding single and double gluon vertex graphs and with g^4 order mass shift terms from the 1-loop diagrams.




FIG. 10. Mass renormalization of order g^4 : cancellation of linear terms in group-5.